

$B \rightarrow X_s l_i^+ l_j^-$ Decays with R-parity ViolationJi-Ho Jang ^{*}, Yeong Gyun Kim [†] and Jae Sik Lee [‡]*Department of Physics, Korea Advanced Institute of Science and Technology
Taejon 305-701, Korea***Abstract**

We derive the upper bounds on certain products of R-parity- and lepton-flavor-violating couplings from $B \rightarrow X_s l_i^+ l_j^-$ decays. These modes of B -meson decays can constrain the product combinations of the couplings with one or more heavy generation indices which are comparable with or stronger than the present bounds. From the studies of the invariant dilepton mass spectrum and the forward backward asymmetry of the emitted leptons we note the possibility of detecting R-parity-violating signals even when the total decay rate due to R-parity violating couplings is comparable with that in the standard model and discriminating two types of R-parity-violating signals. The general expectation of the enhancement of the forward backward asymmetry of the emitted leptons in the minimal supersymmetric standard model with R-parity may be corrupted by R-parity violation.

PACS Number: 11.30.Pb, 11.30.Fs, 13.25.Hw, 13.20.-v

Typeset using REVTeX

^{*}E-mail: jhjang@chep6.kaist.ac.kr

[†]E-mail: ygkim@chep6.kaist.ac.kr

[‡]E-mail: jslee@chep6.kaist.ac.kr

1. In supersymmetric extensions of the standard model, there are gauge invariant interactions which violate the baryon number (B) and the lepton number (L) in general. To prevent occurrences of these B - and L -violating interactions in supersymmetric extensions of the standard model, the additional global symmetry is required. This requirement leads to the consideration of the so called R-parity(R_p). Even though the requirement of R_p conservation makes a theory consistent with present experimental searches, there is no good theoretical justification for this requirement. Therefore the models with explicit R_p -violation have been considered by many authors [1].

To discover the R_p violation in future experiments, we need to know what kinds of couplings are severely constrained by present experimental data. Therefore it is important to constrain the R_p violating couplings from the present data, especially data on the processes forbidden or highly suppressed in the SM. Usually the bounds on the R_p violating couplings with at least two heavy fields are not stronger than those with at most one heavy field.

In this paper, we derive the upper bounds on certain products of R_p and lepton flavor violating couplings from $b \rightarrow sl_i^+ l_j^-$ decays in the minimal supersymmetric standard model (MSSM) with explicit R_p violation. These modes of B -meson decays can constrain the product combinations of the couplings with one or more heavy generation indices.

In the MSSM, the most general R_p -violating superpotential is given by

$$W_{R_p} = \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \lambda''_{ijk} U_i^c D_j^c D_k^c. \quad (1)$$

Here i, j, k are generation indices and we assume that possible bilinear terms $\mu_i L_i H_2$ can be rotated away. L_i and Q_i are the $SU(2)$ -doublet lepton and quark superfields and E_i^c, U_i^c, D_i^c are the singlet superfields respectively. λ_{ijk} and λ''_{ijk} are antisymmetric under the interchange of the first two and the last two generation indices respectively; $\lambda_{ijk} = -\lambda_{jik}$ and $\lambda''_{ijk} = -\lambda''_{ikj}$. So the number of couplings is 45 (9 of the λ type, 27 of the λ' type and 9 of the λ'' type). Among these 45 couplings, 36 couplings are related with the lepton flavor violation.

There are upper bounds on a *single* R_p violating coupling from several different sources [2–5]. Among these, upper bounds from neutrinoless double beta decay [3], ν mass [4] and K^+, t -quark decays [5] are strong. Neutrinoless double beta decay gives $\lambda'_{111} < 3.5 \times 10^{-4}$. The bounds from ν mass are $\lambda_{133} < 3 \times 10^{-3}$ and $\lambda'_{133} < 7 \times 10^{-4}$. From K^+ -meson decays one obtain $\lambda'_{ijk} < 0.012$ for $j = 1$ and 2. These bounds from K^+ -meson decays are basis-dependent [5,6]. Here all masses of scalar partners which mediate the processes are assumed to be 100 GeV. Extensive reviews of the updated limits on a single R_p -violating coupling can be found in [6,7].

There are more stringent bounds on some products of the R_p -violating couplings from the mixings of the neutral K - and B - mesons and rare leptonic decays of the K_L -meson, the muon and the tau [8], B^0 decays into two charged leptons [9], $b\bar{b}$ productions at LEP [10] and muon(ium) conversion, and τ and π^0 decays [11].

In this paper, we assume that the baryon number violating couplings λ'' 's vanish in order to avoid too fast proton decays. Especially in the models with a very light gravitino (G) or axino ($\tilde{a}$), λ'' have to be very small independently of λ' from the proton decay

$p \rightarrow K^+ G$ (or $K^+ \tilde{a}$) ; $\lambda''_{112} < 10^{-15}$ [12]. One can construct a grand unified model which has only lepton number non-conserving trilinear operators in the low energy superpotential when R_p is broken only by bilinear terms of the form $L_i H_2$ [13]. And usually it may be very difficult to discern signals of B -violating interactions above QCD backgrounds [4].

2. The exchange of the sleptons or squarks leads to the four-fermion interactions in the effective lagrangian. Among these four-fermion operators, there are terms relevant for $b \rightarrow sl_i^+ l_j^-$. These effective terms have 2 down-type quarks and 2 charged leptons. From Eq. (1) we obtain

$$\begin{aligned} \mathcal{L}_{R_p}^{eff} = & \sum_{n=1}^3 \frac{2}{m_{\tilde{l}_n}^2} \left[\lambda_{njk}^* \lambda'_{nlm} (\bar{e}_j P_R e_k) (\bar{d}_m P_L d_l) + h.c. \right] \\ & - \sum_{n,r,s=1}^3 \frac{1}{2m_{\tilde{Q}_n}^2} K_{nr} K_{ns}^* \lambda_{jrk}'^* \lambda'_{lsm} (\bar{e}_j \gamma^\mu P_L e_l) (\bar{d}_m \gamma_\mu P_R d_k), \end{aligned} \quad (2)$$

where K is the CKM matrix and we assume the matrices of the soft mass terms are diagonal and the CKM matrix comes from the mixings between down type quarks. From this Lagrangian, one finds that the various semileptonic b decays could appear at tree level; $b \rightarrow qe^+e^-, q\mu^+\mu^-, q\tau^+\tau^-, qe^\pm\mu^\mp, q\mu^\pm\tau^\mp, qe^\pm\tau^\mp$ ($q = d, s$).

At presents, the measurements of the branching ratios of the $b \rightarrow sl_i^+ l_j^-$ processes give the upper bounds (at 90 % C.L.) [14]

$$\begin{aligned} BR(b \rightarrow se^+e^-) & < 5.7 \times 10^{-5}, \\ BR(b \rightarrow s\mu^+\mu^-) & < 5.8 \times 10^{-5}, \\ BR(b \rightarrow se^\pm\mu^\mp) & < 2.2 \times 10^{-5}. \end{aligned} \quad (3)$$

In the SM, the process $b \rightarrow se^\pm\mu^\mp$ is forbidden due to the conservation of each lepton flavor number. On the other hand, the decay $b \rightarrow s\mu^+\mu^-$ ($b \rightarrow se^+e^-$) is dominated by the electroweak penguin and receives small contributions from box diagrams and magnetic penguins. A recent analysis gives $BR(b \rightarrow se^+e^-)_{SM} = (8.4 \pm 2.3) \times 10^{-6}$, $BR(b \rightarrow s\mu^+\mu^-)_{SM} = (5.7 \pm 1.2) \times 10^{-6}$ [16]. The experimental bounds are almost one order of magnitude larger than the standard model expeations. If we neglect the SM contribution, the decay rate of the processes $b \rightarrow sl_i^+ l_j^-$ reads

$$\Gamma(b \rightarrow sl_i^+ l_j^-) = \frac{m_b^5}{6144\pi^3 \tilde{m}^4} \left[4(|\mathcal{A}_{ij}|^2 + |\mathcal{B}_{ij}|^2) + |\mathcal{C}_{ij}|^2 \right] \quad (4)$$

The constants $\mathcal{A}_{ij}$, $\mathcal{B}_{ij}$ and $\mathcal{C}_{ij}$ are given by

$$\mathcal{A}_{ij} = \sum_{n=1}^3 \lambda_{nji}^* \lambda'_{n32},$$

$$\begin{aligned}\mathcal{B}_{ij} &= \sum_{n=1}^3 \lambda_{nij} \lambda_{n23}^*, \\ \mathcal{C}_{ij} &= \sum_{n,p,s=1}^3 K_{np} K_{ns}^* \lambda_{jp3}^* \lambda'_{is2} = \sum_{n=1}^3 \lambda_{jn3}^* \lambda'_{in2}.\end{aligned}\quad (5)$$

Note that we assume the universal soft mass $\tilde{m}$ and ignore the lepton mass. To remove the large uncertainty in the total decay rate associated with the m_b^5 factor, it is convenient to normalize $BR(b \rightarrow sl_i^+ l_j^-)$ to the semileptonic rate $BR(b \rightarrow ce\bar{\nu})$. We then obtain

$$4(|\mathcal{A}_{ij}|^2 + |\mathcal{B}_{ij}|^2) + |\mathcal{C}_{ij}|^2 = \frac{6144\tilde{m}^4 G_F^2 |V_{cb}|^2 f_{\text{ps}}(m_c^2/m_b^2)}{192} \frac{BR(b \rightarrow sl_i^+ l_j^-)}{BR(b \rightarrow ce\bar{\nu})}, \quad (6)$$

where $f_{\text{ps}}(x) = 1 - 8x + 8x^3 - x^4 - 12x^2 \ln x$ is the usual phase-space factor. Inserting into Eq. (6) the semileptonic rate $BR(b \rightarrow ce\bar{\nu}) \approx 10.5\%$, $f_{\text{ps}}(m_c^2/m_b^2) \approx 0.5$, $|V_{cb}| \approx 0.04$, we obtain

$$4(|\mathcal{A}_{ij}|^2 + |\mathcal{B}_{ij}|^2) + |\mathcal{C}_{ij}|^2 = 3.3 \times 10^{-3} \left(\frac{\tilde{m}}{100 \text{ GeV}} \right)^4 BR(b \rightarrow sl_i^+ l_j^-). \quad (7)$$

In the case of $b \rightarrow se^+ e^-$ decay ($i = 1, j = 1$), we obtain

$$4(|\mathcal{A}_{11}|^2 + |\mathcal{B}_{11}|^2) + |\mathcal{C}_{11}|^2 < 1.9 \times 10^{-7}, \quad (8)$$

from the Eq. (7) and the upper limit on the branching ratio, Eq. (3). Under the assumption that only one product combination is not zero, we can get the upper bounds on the following combinations of the $\lambda\lambda'$ - and $\lambda'\lambda'$ -type;

$$\begin{aligned}|\lambda_{n11} \lambda'_{n32}| &< 2.2 \times 10^{-4}, \\ |\lambda_{n11} \lambda'_{n23}| &< 2.2 \times 10^{-4}, \\ |\lambda'_{1n3} \lambda'_{1n2}| &< 4.3 \times 10^{-4}.\end{aligned}\quad (9)$$

For the product combinations of $\lambda\lambda'$ type, we observe that the bounds on $\lambda_{121} \lambda'_{232}$, $\lambda_{131} \lambda'_{332}$, $\lambda_{121} \lambda'_{223}$, $\lambda_{131} \lambda'_{323}$ are stronger than the previous bounds (see Table I).

In a similar way, another upper bounds can be obtained in the case of $b \rightarrow s\mu^+ \mu^-$ and $b \rightarrow se^\pm \mu^\mp$ decays. For the $b \rightarrow s\mu^+ \mu^-$ decay, the bound on $\lambda'_{233} \lambda'_{232}$ is stronger than the previous bounds. And for the $b \rightarrow se^\pm \mu^\mp$ decay, the bounds on $\lambda'_{233} \lambda'_{232}$ and all product combinations of $\lambda\lambda'$ type are stronger than the previous bounds.

In fact, the $B_s \rightarrow l_i^+ l_j^-$ process is described with the same parameters $\mathcal{A}, \mathcal{B}, \mathcal{C}$ and in some cases gives slightly more stringent bounds [9]. But, two things make the decay $b \rightarrow sl_i^+ l_j^-$ more useful than the $B_s \rightarrow l_i^+ l_j^-$ process. One thing is that the contribution of the product combinations of $\lambda'\lambda'$ type is vanishing in the limit of zero lepton mass and so these parameters cannot be constrained in the case of the $B_s \rightarrow l_i^+ l_j^-$ process. The other thing is that the experimental upper limit on the branching ratios exist only in the $B_s \rightarrow \mu^+ \mu^-$ process at present.

3. The structure of R_p effective lagrangian (Eq. (2)) is quite different from that of the SM, it would be interesting to compare the invariant dilepton mass spectrum and the forward backward asymmetry of the emitted leptons in the presence of R_p with those of the SM [16,18] and the minimal MSSM with R_p [19]. The forward backward asymmetry is defined by

$$\frac{d\mathcal{A}(\hat{s})}{d\hat{s}} = \int_0^1 \frac{d^2\mathcal{B}}{d\hat{s}dz} - \int_{-1}^0 \frac{d^2\mathcal{B}}{d\hat{s}dz},$$

where $\hat{s} \equiv s/m_b^2$, s is the invariant mass of the lepton pair, and $z \equiv \cos\theta$ is the angle of l^+ measured with respect to the b -quark direction in the dilepton center of mass system.

Neglecting the masses of the strange quark and leptons, we obtain the R_p invariant dilepton mass spectrum as follows

$$\frac{d\mathcal{B}^{R_p}(b \rightarrow sl_i^+ l_j^-)}{d\hat{s}} = \frac{\mathcal{B}_{sl}(1 - \hat{s})^2 [24(|\mathcal{A}_{ij}|^2 + |\mathcal{B}_{ij}|^2)\hat{s} + |\mathcal{C}_{ij}|^2(1 + 2\hat{s})]}{16f_{ps}(m_c^2/m_b^2)G_F^2\tilde{m}^4|V_{cb}|^2}, \quad (10)$$

where $\mathcal{B}_{sl}$ is the semileptonic rate $BR(b \rightarrow ce\bar{\nu})$. We find that there are *no* interferences between contributions of the SM and the R_p -violating model under consideration. So, the total invariant dilepton mass spectrum is given by the direct sum of the SM contributions and R_p -violating contributions. In Fig. 1 (a), we show the normalized invariant dilepton mass spectrum $\frac{d\mathcal{B}}{d\hat{s}} \equiv \frac{1}{\mathcal{B}} \frac{d\mathcal{B}}{d\hat{s}}$. The solid line denotes the typical SM prediction [16,18] and dashed and dotted lines the R_p -violating contributions to the invariant dilepton mass spectrum. Since the structure of the effective lagrangian corresponding to $\lambda\lambda'$ -type R_p violation is different from that of $\lambda'\lambda'$ -type one, the $\hat{s}$ dependences of the invariant dilepton mass spectrum due to these two types of R_p violation are quite different from each other. The $\hat{s}$ dependence of $\lambda'\lambda'$ -type R_p violation is nearly the same as that of the SM and this type of R_p violation enhances the decay rate on the low $\hat{s}$ region. But the behavior of the contribution from $\lambda\lambda'$ -type R_p violation is different from the SM and this type of R_p violation enhances the invariant dilepton mass spectrum in the region around $\hat{s} = 1/3$. Therefore, from the invariant dilepton mass spectrum one can detect the R_p violating signals even if the magnitude of R_p violating coupling is comparable with that of the SM and it is possible to discriminate two types of R_p violating signals, $\lambda\lambda'$ - and $\lambda'\lambda'$ -type.

We also obtain R_p forward backward asymmetry of the emitted leptons

$$\frac{d\mathcal{A}^{R_p}(b \rightarrow sl_i^+ l_j^-)}{d\hat{s}} = \frac{-3\mathcal{B}_{sl}|\mathcal{C}_{ij}|^2(1 - \hat{s})^2\hat{s}}{32f_{ps}(m_c^2/m_b^2)G_F^2\tilde{m}^4|V_{cb}|^2}. \quad (11)$$

Let us note that sign of this asymmetry is negative. In Fig. 1 (b), we show the normalized forward backward asymmetry of the emitted leptons $\frac{d\bar{\mathcal{A}}}{d\hat{s}} \equiv \frac{1}{\mathcal{B}} \frac{d\mathcal{A}}{d\hat{s}}$. There is no contribution to the asymmetry from the $\lambda\lambda'$ -type R_p violation. But there is *negative* contribution from $\lambda'\lambda'$ -type R_p violation which can compensate the SM asymmetry. Since there are no interferences between contributions of the SM and the R_p -violating model under consideration, the total asymmetry $\frac{d\bar{\mathcal{A}}}{d\hat{s}}$ is given by

$$\left(\frac{d\tilde{\mathcal{A}}}{d\hat{s}}\right) \equiv \frac{\left(\frac{d\mathcal{A}}{d\hat{s}}\right)^{\text{SM}} + \left(\frac{d\mathcal{A}}{d\hat{s}}\right)^{R_p}}{\left(\frac{d\mathcal{B}}{d\hat{s}}\right)^{\text{SM}} + \left(\frac{d\mathcal{B}}{d\hat{s}}\right)^{R_p}} = \frac{\left(\frac{d\bar{\mathcal{A}}}{d\hat{s}}\right)^{\text{SM}} + \xi_{\lambda\lambda'} \left(\frac{d\bar{\mathcal{A}}}{d\hat{s}}\right)^{R_p}_{\lambda\lambda'} + \xi_{\lambda'\lambda'} \left(\frac{d\bar{\mathcal{A}}}{d\hat{s}}\right)^{R_p}_{\lambda'\lambda'}}{\left(\frac{d\bar{\mathcal{B}}}{d\hat{s}}\right)^{\text{SM}} + \xi_{\lambda\lambda'} \left(\frac{d\bar{\mathcal{B}}}{d\hat{s}}\right)^{R_p}_{\lambda\lambda'} + \xi_{\lambda'\lambda'} \left(\frac{d\bar{\mathcal{B}}}{d\hat{s}}\right)^{R_p}_{\lambda'\lambda'}}, \quad (12)$$

where $\xi_{(\lambda\lambda',\lambda'\lambda')} \equiv \mathcal{B}_{(\lambda\lambda',\lambda'\lambda')}^{R_p}/\mathcal{B}^{\text{SM}}$ and $\left(\frac{d\bar{\mathcal{A}}}{d\hat{s}}\right)^{R_p}_{\lambda\lambda'} = 0$. In the case $\xi_{\lambda\lambda'} \gg \xi_{\lambda'\lambda'}$, the SM asymmetry will be diluted by the factor of $\sim 1/(1 + \xi_{\lambda\lambda'})$. In the case $\xi_{\lambda\lambda'} \ll \xi_{\lambda'\lambda'}$, even the sign of the asymmetry could be different from the prediction of the SM depending on the size of $\xi_{\lambda'\lambda'}$. For illustration, we show the above forward backward asymmetry with $\xi_{\lambda\lambda'} = 1$ or $\xi_{\lambda'\lambda'} = 1$ in Fig. 2. So, also from the forward backward asymmetry of the emitted leptons one can detect the R_p violating signals even if the magnitude of R_p violating coupling is comparable with that of the SM and it is possible to discriminate two types of R_p violating signals, $\lambda\lambda'$ - and $\lambda'\lambda'$ -type.

From the studies of the process $b \rightarrow sl^+l^-$ in the minimal MSSM with R_p [19], the SUSY signal may appear as the enhancement of this asymmetry by more than 100 % relative to SM expectations. But, in the presence of R_p violation this minimal MSSM enhancement may be corrupted ¹.

4. To conclude, we have derived the more stringent upper bounds on certain products of R_p - and lepton-flavor-violating couplings from the recent measurements of $b \rightarrow sl_i^+l_j^-$ decay rates at CLEO. From the studies of the invariant dilepton mass spectrum and the forward backward asymmetry of the emitted leptons, we note the possibility of detecting R_p -violating signals even when the total decay rate due to R_p -violating couplings is comparable with that in the SM and discriminating two types of R_p -violating signals, $\lambda\lambda'$ - and $\lambda'\lambda'$ -type. The general expectation of the enhancement of the forward backward asymmetry of the emitted leptons in the minimal MSSM with R_p may be corrupted by R-parity violation.

ACKNOWLEDGEMENTS

We thank P. Ko and K. Choi for helpful discussions. This work was supported in part by the KRF postdoctoral program (Y.G.K.) and in part by the KAIST Basic Science Research Program (J.S.L.).

¹In fact, the precise definition of the asymmetry in the first reference of [19] is different from ours. But, our conclusion is not changed by this difference.

REFERENCES

- [1] J. Ellis et al., Phys. Lett. **150B**, 142 (1985); G. G. Ross and J. W. F. Valle, Phys. Lett. **151B**, 375 (1985); S. Dawson, Nucl. Phys. **B261**, 297 (1985); S. Dimopoulos and L. Hall, Phys. Lett. **207B**, 210 (1987).
- [2] V. Barger, G. F. Giudice and T. Han, Phys. Rev. **D40**, 2987 (1989).
- [3] R. N. Mohapatra, Phys. Rev. **D34**, 3457 (1986); M. Hirsch, H. V. Klapdor-Kleingrothaus and S. G. Kovalenko, Phys. Rev. Lett. **75**, 17 (1995).
- [4] R. M. Godbole, P. Roy and X. Tata, Nucl. Phys. **B401**, 67 (1993).
- [5] K. Agashe, M. Graesser, Phys. Rev. **D54**, 4445 (1996).
- [6] G. Bhattacharyya, Nucl. Phys. Proc. Suppl. **52A**, 83 (1997), G. Bhattacharyya, hep-ph/9709395.
- [7] M. Chaichian and K. Huitu, Phys. Lett. **384B**, 157 (1996).
- [8] D. Choudhury and P. Roy, Phys. Lett. **378B**, 153 (1996).
- [9] J. Jang, J. K. Kim and J. S. Lee, Phys. Rev. **D55**, 7296 (1997).
- [10] J. Erler, J. L. Feng and N. Polonsky, Phys. Rev. Lett. **78**, 3063 (1997)
- [11] J. E. Kim, P. Ko and D. Lee, Phys. Rev. **D56**, 100 (1997).
- [12] K. Choi, E. J. Chun and J. S. Lee, Phys. Rev. **D55**, 3924 (1997)
- [13] L. J. Hall and M. Suzuki, Nucl. Phys. **B231**, 419 (1984); F. Vissani, Nucl. Phys. Proc. Suppl. **52A**, 94 (1997).
- [14] CLEO Collaboration, hep-ex/9710003
- [15] Ref. [108] in A. J. Buras, hep-ph/9610461
- [16] A. Ali, G. Hiller, L. T. Handoko and T. Morozumi, Phys. Rev. **D55**, 4105 (1997).
- [17] J. Jang, Y. G. Kim and J. S. Lee, Phys. Lett. **408B**, 367 (1997)
- [18] M. Misiak, Nucl. Phys. **B393**, 23 (1993), Erratum, Nucl. Phys. **B439**, 461 (1995), A. J. Buras and M. Münz, Phys. Rev. **D52**, 186 (1995), A. J. Buras R. Fleisher, hep-ph/9704376.
- [19] P. Cho, M. Misiak and D. Wyler, Phys. Rev. **D54**, 3329 (1996), A. Masiero and L. Silvestrini, hep-ph/9709244.

TABLES

TABLE I. Upper bounds on the magnitudes of products of couplings derived from $b \rightarrow sl_i^+ l_j^-$.

Decay Mode	Combinations Constrained	Upper bound	Previous bound
$b \rightarrow se^+ e^-$	$\lambda_{121} \lambda'_{232}$	2.2×10^{-4}	$8.0 \times 10^{-3} \text{ }^a$
	$\lambda_{131} \lambda'_{332}$	2.2×10^{-4}	2.9×10^{-2}
	$\lambda_{121} \lambda'_{223}$	2.2×10^{-4}	$6.0 \times 10^{-4} \text{ }^b$, $(9.0 \times 10^{-3} \text{ }^c)$
	$\lambda_{131} \lambda'_{323}$	2.2×10^{-4}	$7.2 \times 10^{-4} \text{ }^b$, $(1.2 \times 10^{-2} \text{ }^c)$
	$\lambda'_{113} \lambda'_{112}$	4.3×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(4.0 \times 10^{-4} \text{ }^c)$
	$\lambda'_{123} \lambda'_{122}$	4.3×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(4.0 \times 10^{-3} \text{ }^c)$
	$\lambda'_{133} \lambda'_{132}$	4.3×10^{-4}	$1.1 \times 10^{-4} \text{ }^a$
$b \rightarrow s\mu^+ \mu^-$	$\lambda_{122} \lambda'_{132}$	2.2×10^{-4}	$5.5 \times 10^{-5} \text{ }^d$
	$\lambda_{232} \lambda'_{332}$	2.2×10^{-4}	$5.5 \times 10^{-5} \text{ }^d$
	$\lambda_{122} \lambda'_{123}$	2.2×10^{-4}	$5.5 \times 10^{-5} \text{ }^d$
	$\lambda_{232} \lambda'_{323}$	2.2×10^{-4}	$5.5 \times 10^{-5} \text{ }^d$
	$\lambda'_{213} \lambda'_{212}$	4.4×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(8.1 \times 10^{-3} \text{ }^c)$
	$\lambda'_{223} \lambda'_{222}$	4.4×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(3.2 \times 10^{-2} \text{ }^c)$
	$\lambda'_{233} \lambda'_{232}$	4.4×10^{-4}	$2.5 \times 10^{-2} \text{ }^a$
$b \rightarrow se^\pm \mu^\mp$	$\lambda_{122} \lambda'_{232}$	1.4×10^{-4}	$1.8 \times 10^{-2} \text{ }^a$
	$\lambda_{132} \lambda'_{332}$	1.4×10^{-4}	2.9×10^{-2}
	$\lambda_{121} \lambda'_{132}$	1.4×10^{-4}	$8.0 \times 10^{-3} \text{ }^a$
	$\lambda_{231} \lambda'_{332}$	1.4×10^{-4}	2.9×10^{-2}
	$\lambda_{122} \lambda'_{223}$	1.4×10^{-4}	$6.0 \times 10^{-4} \text{ }^b$, $(9.0 \times 10^{-3} \text{ }^c)$
	$\lambda_{132} \lambda'_{323}$	1.4×10^{-4}	$7.2 \times 10^{-4} \text{ }^b$, $(1.2 \times 10^{-2} \text{ }^c)$
	$\lambda_{121} \lambda'_{123}$	1.4×10^{-4}	$6.0 \times 10^{-4} \text{ }^b$, $(1.0 \times 10^{-2} \text{ }^c)$
	$\lambda_{231} \lambda'_{323}$	1.4×10^{-4}	$7.2 \times 10^{-4} \text{ }^b$, $(1.2 \times 10^{-3} \text{ }^c)$
	$\lambda'_{113} \lambda'_{212}$	2.7×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(1.8 \times 10^{-3} \text{ }^c)$
	$\lambda'_{123} \lambda'_{222}$	2.7×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(3.6 \times 10^{-2} \text{ }^c)$
	$\lambda'_{133} \lambda'_{232}$	2.7×10^{-4}	$1.1 \times 10^{-4} \text{ }^a$
	$\lambda'_{213} \lambda'_{112}$	2.7×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(1.8 \times 10^{-3} \text{ }^c)$
	$\lambda'_{223} \lambda'_{122}$	2.7×10^{-4}	$1.4 \times 10^{-4} \text{ }^b$, $(3.6 \times 10^{-3} \text{ }^c)$
	$\lambda'_{233} \lambda'_{132}$	2.7×10^{-4}	$2.5 \times 10^{-2} \text{ }^a$

a: Bounds from $B \rightarrow X_c l \bar{\nu}$ [17]. b: Considering the bounds from $K \rightarrow \pi \nu \bar{\nu}$ [5]. c: Ignoring the bounds from $K \rightarrow \pi \nu \bar{\nu}$. d: Bounds from $B_s \rightarrow \mu^+ \mu^-$ [9]. Others : See the second reference of [6].

FIGURES




Fig. 1

FIG. 1. Plots of normalized invariant dilepton mass spectrum (a) and forward backward asymmetry of the emitted leptons (b). The solid line denotes the typical SM prediction [16,18] and dashed and dotted lines the R_p -violating contributions, Eq.(10) and Eq. (11).




FIG. 2. Plots of total forward backward asymmetry of the emitted leptons, $\left(\frac{d\bar{A}}{d\hat{s}}\right)$. The solid line denotes the typical SM prediction [16,18] and dashed and dotted lines the R_p -violating contributions, Eq.(12).